

## Exploring the bioefficacy of Endophytic Bacteria against Important Plant Pathogens

**Seweta Srivastava<sup>1\*</sup>., Aspak<sup>1.</sup>., Meenakshi Rana<sup>1.</sup>., Kanuri Komala Siva Katyayani<sup>2.</sup>., Dipshikha Kaushik<sup>2.</sup>., Rajeev Kumar<sup>3.</sup>., Manash Shukla<sup>1.</sup>., Shubham Kumar<sup>1.</sup>., Raghavendra Reddy Manda<sup>4</sup> and Vinit Pratap Singh<sup>5</sup>**

### ABSTRACT

The biological management of plant diseases has developed into a separate scientific and technological discipline, and in recent years, this change has happened quickly. A form of bacterium known as a bacterial endophyte may colonize any portion of a plant without causing any symptoms or harm to the host plant. Endophytic bacteria have been discovered by several researchers, and there is growing evidence that they can stop a variety of plant diseases from growing and functioning. Endophytes have a variety of benefits including growth-increasing and disease-hampering properties. Researchers' interest in this field is growing as a result of its potentially to be utilized as an alternative to synthetic fungicides. This review's main objectives are to chart the development of endophytic bacterial research and give scientists access to current knowledge that will spur further investigation. Endophytic bacteria are employed to control plant diseases including wilt, rot and post-harvest damage, as well as nematode infestation. Endophytic bacteria are also used to control nematodes and postharvest diseases. With an emphasis on endophytic bacteria, this review explains the diverse mechanisms of bacterial endophytes to shield the plant from biotic infection.

**Keywords:** Antibiotics, Bacteria, Endophytes, Management, Phytopathogens

**MS History:** 24.03.2023(Received)-30.05.2023(Revised)- 10.06.2023 (Accepted)

**Citation:** Seweta Srivastava, Aspak., Meenakshi Rana., Kanuri Komala Siva Katyayani., Dipshikha Kaushik., Rajeev Kumar., Manash Shukla., Shubham Kumar., Raghavendra Reddy Manda and Vinit Pratap Singh. 2023. Exploring the bioefficacy of Endophytic Bacteria against Important Plant Pathogens. *Journal of Biopesticides*, 16(1):79-99. DOI:10.57182/jbiopestic.16.1.79-99

### INTRODUCTION

Plant diseases create tremendous biotic stress in plants, causing farmers to lose a lot of money and tainting food by creating toxins while it is kept. Farmers' purposeful determination to combat illness resulted in the development of a variety of pesticidal molecule, the use of which destroys the environment and eventually, harms human health. Plant health management has gotten more difficult as certain plant diseases have developed resistance to these treatments (Dun-chun *et al.*, 2016). Bio-control of plant diseases has become more important in addressing these concerns. Plant growth-promoting rhizobacteria (PGPR) have long

been investigated by many scientists and rhizosphere treatments for biocontrol have mostly focused on them. Due to the expanding range of ways that microorganisms may be used to boost plant development and lower disease-causing pathogens, researchers have lately turned their attention to those that colonize interior tissues with laser beams (Saeed *et al.*, 2021). Researchers have recently focused a lot of emphasis on the function of bacterial endophytes among these microorganisms in plant disease management. Endophytic bacteria were defined by Wilson (1995) as prokaryotes that seek to colonize the vascular tissues without causing any damage to the

host plant. Endophytes are "endo-symbionts" that live inside plant tissues without causing injury or illness and may be discovered using aseptic procedures, according to researchers. Previous studies showed the beneficial relations between plants and microorganisms and scientists believed that fungi that weren't often recognized for causing illnesses in agricultural plants had the power of microbial endophytes (Clay, 1988). The seeds of horticultural as well as agricultural crops might be used to isolate bacterial species (Kirchhof *et al.*, 1997).

According to studies, endophytic bacteria can be found in plant parts. When describing the habitat of endophytes, Andrews (1992) stated that, unlike microorganisms dwelling in and above the rhizosphere, endophytes may exist in a fully isolated environment. Endophytic bacteria, according to Arnold and Lutzoni (2007), may reside in the rhizosphere, twig, leaves, petals, seeds and fruits of agricultural plants.

Endophytes have a variety of benefits, according to a growing body of literature. Kang *et al.*, (2007) described endophytes growth-increasing properties, whereas Senthilkumar *et al.*, (2007) performed endophytes' disease-hampering properties. Bakker *et al.*, (2007) investigated the work of endophytes in strengthening crop defense mechanisms against various plant disease. Endophytes have been shown to generate anti-herbivory compounds as well as catalyze biological nitrogen fixation in plants (Martínez *et al.*, 2003) and improve their mineral absorption (Malinowski *et al.*, 2000). Backman *et al.*, (1997) conferred specific bacteria colonizing a specific crop species, changing populations as seasons change, the order in which they colonized and their capability to mobilize within cells and encourage systemic resistance as endophytes as antimicrobials against multiple plant diseases.

### **Endophytes**

A quick description of 'Endophytes' is provided here to help you comprehend the subsequent sections of the review. Endophytes are microorganisms that be inherent asymptotically in the plant for at least a portion of their lifespan (Solis *et*

*al.*, 2016). Endophytes thrive within their hosts intracellularly, systemically or locally without creating apparent infection or disease signs (Schulz *et al.*, 2015). According to Busby *et al.*, (2016), endophytism is characterized by "inconspicuous infections, diseased host tissues that are at least temporally symptomless and demonstrated microbial colonization inside host tissues". All plants are thought to have endophytes, and the biodiversity of these microorganisms relies on a range of factors, including the type of host plant, plant canopy, nutrient availability, the adequacy of the local environment and interactions between bacteria and fungi that are carried by the soil (Yan *et al.*, 2015). Endophytes are potential biocontrol agents because they can change interactions with infections and pests. An endophyte called *Acremonium alternatum* boosts tomato resistance to the powdery mildew disease *Leveillula taurica* and shields beans from the moth *Plutella xylostella*. An isolated fungal endophyte from cotton plants called Phomopsis sp. prevented caterpillar herbivory on cotton plants. Sometimes an endophyte species can act as a biocontrol agent, and other times it might promote the growth of the host plant, which has additional benefits. Neotyphodium species promote host plant growth, fitness and stress tolerance while safeguarding it against infections and pests (Solis *et al.*, 2016). Furthermore, pathogenic *Sclerotium rolfsii* was decreased and sunflower biomass output was boosted by endophytic *Penicillium citrinum* and *Aspergillus terreus* (Harman *et al.*, 2021). How endophytes minimize diseases and pests is the next important question. We will explore how endophytes maintain their interaction with their hosts before diving into several biocontrol techniques.

### **Interaction between plants and endophytes**

The concept of "balanced antagonism" between endophytes and their host explains why they colonize without exhibiting any symptoms (Schulz *et al.*, 2015). Fungal virulence factors will be totally overcome by plant defence systems,

preventing the fungus from colonising plant tissues. If fungal virulence elements could interfere with plant defence systems, a plant-pathogen connection would result in plant disease (Suryanarayanan *et al.*, 2016).

When they are impacted by internal or external conditions that make them express pathogenic factors, certain endophytes turn into pathogens (Kusari *et al.*, 2012). *Colletotrichum magna* strains that are pathogenic and endophytic have been demonstrated to transform their life styles by interfering with certain genetic loci or closely related genes that cause anthracnose disease in cucurbitaceous crop (Rai and Agarkar, 2016). A non-pathogenic mutant strain of *Colletotrichum magna* (Path-1) produced from a pathogenic strain (CmL2.5) colonizes the roots and stems of cucurbit plants asymptotically and inhibits the virulent form of the fungus, according to experiments (Rai and Agarkar, 2016). High humidity or a shortage of nutrients may be to blame for this frequent occurrence of *Colletotrichum* switching lifestyles, which alters the host's vulnerability in the presence of natural circumstances (Fisher and Petrini, 1992; Rai and Agarkar, 2016).

Some endophytes produce small quantities of antifungal and antibacterial chemicals to prevent competitors (both pathogenic and endophytic bacteria and fungi) and maintain a competitive balance (Suryanarayanan *et al.*, 2016). The insecticidal metabolite rugulosin generated by endophytic *Phialocephala* species from *Picea glauca* (white spruce) poisons *Choristoneura fumiferana* (spruce bud worm). Secondary metabolites regulate the antagonistic connections between competitors, plant hosts, and endophytes (Hashem *et al.*, 2023). Estrada *et al.*, (2012) found that endophytic *Fusarium verticillioides* in maize might lower pathogenic *Ustilago maydis* aggressiveness while simultaneously destroying protective systems.

The compounds in the plant are effective against *U. maydis*. Pathogen reduction may also come through multipartite healthy relations between endophytes, competitors and host plants.

Secondary metabolites will impair their ability to develop and survive (Suryanarayanan *et al.*, 2016). In conclusion, interactions between plants and endophytes are complex and control the balance of host defence, fungal virulence and secondary metabolites.

#### **Metabolites and activities of endophytes**


The potentiality of microbial endophytes to yield a variety of crucial compounds for pharmacology, including antiviral, antifungal, antibacterial, antitumor and anticancer medications, is well documented. Several endophytes can produce plant hormones and growth factors (Kandel *et al.*, 2017; Chaudhary *et al.*, 2022). Abiotic stress tolerance, siderophores, nematocidal, insecticidal and agricultural chemicals are some of their other potential products. A variety of extracellular enzymes, including the phosphatase enzyme, which transforms insoluble phosphate into soluble phosphates for easier digestion by plants, have been shown to be secreted by endophytes (Sharma *et al.*, 2021). Endophytes create chemicals that can be employed in the production of biofuels and the degradation of sophisticated organic and inorganic pollutants that are produced during industrial operations (Burrage and Jeon, 2021). The advantages of endophytes are listed below, along with some prospective uses for them in various industries.

#### **Endophytes potential in agriculture**

Endophytes, according to published studies, are a good source of metabolites and desirable functionalities that might benefit an organic agricultural system. Some endophytes might be employed as bio-pesticides against plant pathogens because of their antibacterial, nematocidal and insecticidal capabilities.

#### **Biopesticidal properties of Endophytes**

A systemic weed commensal fungal endophyte *Epichloe typhina* releases mycotoxic properties in extracts of *Phleum pratense*, a perennial grass native to much of Europe. Bacteria generated chitinase, which is known to dissolve chitin polymers, which are a key component of a fungal cell wall. *Bacillus cereus* strain was recognized as



**Fig.1.** Endophytes and their diverse properties (Source: Unpublished photographs from the authors)

bacterial endophyte, was previously perform a defense mechanism against *Rhizoctonia solani* (Pleban *et al.*, 1997). A strain of *Neotyphodium sp.* (AR601) that produces substantial amounts of alkaloids such as loline and ovaline and is injected into the turf tall fescue cultivar 'Jackal' has shown

bird deterring capacity (Pennell, 2010). By generating pathogenesis-related proteins, some endophytes have been confirmed to reliably produce effective resistance in plants against common phytopathogens. Fungal endophytes isolated from the tree leaves were shown to produce chitinase and chitosanase, which may help

Table 1. Mechanism involved in the mode of action of bacterial endophytes

Broad mode of action	Mechanism involved	References
Root colonization through competition	Various growth stages, the capability to adhere to roots and circulate around without inhibition, and the efficient utilization of the organic acids released from root exudates, the generation of a range of chemicals, together with amino acids, and the type III secretion system are all characteristics of this species.	Lugtenberg and Kamilova, 2009
Antibiosis and antibiotics suppressing pathogens	Pharmaceuticals such as phenazines, pyoluteorin, pyrrolnitrin, and the volatile HCN are produced.	Pierson and Pierson, 2010; Dandurishvili <i>et al.</i> , 2011; Henry <i>et al.</i> , 2011; Savadogo <i>et al.</i> , 2011; Ramkumar <i>et al.</i> , 2013; Zhang <i>et al.</i> , 2013; Torres <i>et al.</i> , 2016
	There is the production of D-gluconic acid, 2-hexyl-5-propyl resorcinol, and the volatiles 2,3-butanediol, 6-pentyl—pyrone, and DMDS.	
	Lipopeptides with disease-controlling abilities include surfactin, fengycin, polymyxin, bacitracin, and the iturin group.	
	Pyrrolnitrin, pyrrologlucinol, phenols, and volatile organic compounds such benzothiazole, pyrazine (2,5-dimethyl), and phenolic derivatives are produced.	
Signal interference	Exo-enzyme synthesis requires the deactivation of AHL molecules.	Dandurishvili <i>et al.</i> , 2011
Ferric iron ion competition	Siderophores are synthesized in order to trap ferric ion.	Whipps, 2001
Competition for nutrients and niches (CNN)	CNN follows the same method as competitive root colonization.	Malfanova, 2013
Detoxification and degradation of virulence factors	Fusaric acid detoxifies toxins released by pathogens.	Uroz <i>et al.</i> , 2003
	By destroying autoinducer signals, which prevent the expression of several virulence genes, the ability to sense quorum is achieved.	
	Resistance produced by salicylic acid, c-LPs, pyocyanins, siderophores, and other substances	

host plants defend against many plant pathogens by activating host defenses and enhancing resistance (Zheng *et al.*, 2017).

#### Antimicrobial properties of endophytes

Some endophyte species have been found to form antimicrobial compounds (Jha *et al.*, 2023). For

their antibacterial properties, endophytic microbes from plants have also been taken into consideration (Wang *et al.*, 2019; Xu *et al.*, 2020). Phomopsichalasin was extracted from *Phomopsis* sp., isolate no. MF6031, which was attained from the twigs of *Salix gracilostyla* var. *melanostachys* was shown to have antibacterial action against *Bacillus subtilis*, *Salmonella gallinarium* and *Staphylococcus aureus* as well as antagonistic activity against *Candida tropicalis* (Horn *et al.*, 1995). In one more investigation, a *Colletotrichum* spp. isolated from internal stem cells of *Artemisia annua* L. was found to exhibit antifungal, antibacterial and fungistatic activities (Lu *et al.*, 2000).

#### **Direct inhibition on plant pathogens**

Several recent research has initiated that endophytes may defend the host plants from diseases or may decrease the destruction triggered by pathogenic microorganisms (Ganley *et al.*, 2008; Meja *et al.*, 2008). Despite the fact that certain research suggests potential endophyte mechanisms for limiting pathogen damage, our understanding of the exact control of endophyte, pathogen and plant is still in its infancy. In this part, we will talk about the processes as direct effects, indirect effects by increasing plant defence and ecological effects. During direct influence, endophytes actively conquer plant diseases by generating antibiotics and lytic enzymes (Fadiji and Babalola, 2020). Conversely, direct interactions amongst bacterial endophytes and biotic plant diseases can be challenging and hostile depending on the species involved (Afzal *et al.* 2019).

#### **Indirect effects of on host plant resistance**

In reaction to severe environmental circumstances such as drought, cold, salt stress or during biotic infections, plants generate a number of defence mechanisms. In response to diverse stimuli, rapid structural and biochemical changes occur, such as cellular necrosis, hypersensitive response and phytoalexin synthesis. Over time, two forms of innate resistance develop to withstand pathogen infestation: non-specific (generic) resistance and

particular resistance (Kira'ly *et al.*, 2007). The previous one is efficient compared to a wide range of pathogenic microbial species, whereas the latter can tolerate infection by a few pathogenic strains. In fact, resistance improvement and secondary metabolite synthesis boost plant defence against endophytes.

#### **Plant Disease Management**

Endophytic bacteria have arisen as an attractive, promising and ecologically friendly biological control technique because they can efficaciously decrease biotic disease incidence and severity by blocking the vascular development of the target pathogen (Constantin *et al.* 2019; de Lamo *et al.* 2018). These endophytes infiltrate plant portions without causing harm. On a variety of hosts, they either directly or indirectly promote plant growth and/or also act as biocontrol agents by inducing resistance (Constantin *et al.* 2019).

#### **Wilt-Causing Pathogens by Bacterial Endophytes**

Wilt is a widespread disease caused by fungal and bacterial strains that can cause major financial losses for farmers. *Fusarium* and *Verticillium* are two significant fungal species that produce wilt, and they are difficult to treat since they are soilborne diseases. The pathogenic agent's soilborne origin and capability to infiltrate the vascular system of infected plants, as well as the rise of new and vigorous pathogen physiological races, make disease treatment difficult. Chemical wilt treatments are generally unsuccessful due to the pathogen's extensive host range and ability to live in soil for lengthy periods of time. As a result, biological wilt management has become more significant, encouraging many scientists to do research on discovering appropriate endophytic bacteria to control wilt infections. Endophytic microorganisms may constitute a potentially appealing and ecologically safe option for wilt pathogen biocontrol because endophytes may better restrict disease occurrence and severity by inhibiting systemic fungal progress (Aydi-Ben-Abdallah *et al.*, 2020). Endophytic bacteria by their diverse mode of action have been revealed in

a quantity of studies to check the growth of wilt-producing pathogens (Table 2).

### Managing Root Rot by Endophytic Bacteria

Pathogens that cause root rot are particularly challenging to control because they may persist in the plant debris/soil up to many years until the environmental conditions are conducive for them and a susceptible host plant can be produced (Conner *et al.*, 2014). The primary method for controlling these infections still involves the use of agrochemicals, but this method has repeatedly led to the emergence of resistance and had a negative impact on the environment. Although frequently employed to address root rots, seed coating with fungicides has had little impact on the pathogens' control (Xu and Kim, 2014). Endophytic bacteria have been praised to manage root rot pathogens because they share a niche with the disease, secrete antifungal metabolites, and aid flora in acquiring nutrients and preparing for plant defence (Muthukumar and Bhaskaran, 2007). Root tissues are colonized by endophytic bacteria, which can defend their host plants from invasion by soil-borne pathogens (Mercado-Blanco *et al.*, 2004; Rybakova *et al.*, 2016) because endophytes are initially seen in root hairs during the initial stages of their colonization, and afterwards move in the root cortex (Prieto *et al.*, 2011; Castanheira *et al.*, 2017; Rangjaroen *et al.*, 2017). Plants benefit from endophytic bacteria invading interior plant tissue in many different ways, with the production of plant growth regulators, osmo-protectants (Beneduzi *et al.*, 2012), exopolysaccharides (Berg *et al.*, 2013), antifungal metabolites (Gond *et al.*, 2015) and regulation of plant physio-biochemical components (Hashem *et al.*, 2016). Regardless of how crucial the endophyte-plant interaction is, little is known about how pathogens, endophytes, and legumes interact in adverse environmental conditions. Management of various rot causing pathogens by endophytic bacteria is summarized in Table 3 mentioned below.

However, only a few endophytic biological control agents have been approved for practice in sustainable agriculture and are currently commercially accessible. This calls for greater

research on the exploration and expansion of biocontrol organisms, particularly the utilization of endophytes.

### Bacterial Endophytes for storage pest

Latest findings have documented the antagonistic behaviors of a wide variety of bacterial endophytes that are found on the outer most layer of fruits and vegetables. On the surface of the fruit, several bacterial species and actinomycetes can influence the development of postharvest diseases (Huang *et al.*, 2021). Three primary bacterial phyla—*Proteobacteria*, *Actinobacteria* and *Bacteroidetes*—dominate the various microbial communities found within or on the host plant surface (Hacquard *et al.*, 2015). The most common biocontrol bacteria discovered on fruit surfaces include *Bacillus* spp., *Burkholderia*, *Citrobacter*, *Pseudomonas* and *Paenibacillus*, (Huang *et al.*, 2021). By displaying antibiosis, *Pantoea dispersa* prevented sweet potato from developing black rot (Jiang *et al.*, 2019). *Streptomyces* species, a Gram-positive bacterium was recently discovered to be able to stop the infection caused by various bacteria and fungi, including *Burkholderia glumae*, a bacterial rice pathogen (Degrassi and Carpentieri-Pipolo 2020).

Notably important tasks are screening microbial antagonists against diverse phytopathogens (Kumari *et al.*, 2022). For BCA screening, bacterial strains that may produce antibiotic or volatile chemicals as well as enzymes that can disrupt or lessen the pathogen virulence factors are favored (Zimand *et al.*, 1996; Kapat *et al.*, 1998; Kumari *et al.*, 2022). Table 4 enlists the endophyte-produced bioactive compounds that may be employed to combat biotic infections after harvest.

### Endophytic in nematodes management

Since the middle of the 1990s, bacterial endophytes have been revealed to be antagonistic to phytopathogenic nematode (Hallmann *et al.*, 1997; Siddiqui and Mahmood, 1999; Bhat *et al.*, 2023). Plant pathogens are opposed by the greater number of Gram-negative endophytic bacteria and by only few species of Gram-positive bacterial

endophyte (Kobayashi and Palumbo, 2000). Gram-negative endophytes include *Burkholderia cepacia*, *P. fluorescens* and *Agrobacterium radiobacter*, whereas Gram-positive endophytes include *Bacillus* spp. *Acromobacter*, *Acinetobacter*, *Agrobacterium*, *Bacillus*,

*Brevibacterium*, *Microbacterium*, *Pseudomonas*, *Xanthomonas* and other species have also been discovered to have the capacity to suppress phytopathogenic nematodes (Yadav *et al.*, 2017; Harni *et al.*, 2023).

Table 2. Role of bacterial endophytes in wilt disease management

Sr No.	Pathogens causing wilt	Endophytic bacteria have been shown to reduce wilt incidence	Mode of action	References
1	<i>Verticillium dahliae</i> F. <i>oxysporum</i> f. Sp. <i>lycopersici</i> F. <i>oxysporum</i> f. Sp. <i>radicislycopersici</i>	<i>Pseudomonas</i> sp. strain PsJN <i>P. fluorescens</i> WCS417r <i>B. pumilus</i> SE-34 <i>Bacillus amyloliquefaciens</i> BO7 <i>B. amyloliquefaciens</i> RWL-1	Endophytic bacteria colonize tomato plants and thicken their cortical cell walls as structural barrier. Siderophores and plant defence hormones like jasmonic acid, and salicylic acid are generated, enhancing ISR.	Vitullo <i>et al.</i> , 2012; Shahzad <i>et al.</i> , 2017
2	<i>F. oxysporum</i> f. Sp. <i>vasinfectum</i> <i>Verticillium dahliae</i>	<i>Aureobacterium saperdae</i> , <i>Bacillus pumilus</i> , <i>Burkholderia solanacearum</i> , <i>Phyllobacterium rubiacearum</i> , <i>Pseudomonas putida</i> , <i>Bacillus subtilis</i> KDRE01, <i>Bacillus megaterium</i> KDRE25	Antibiosis is performed by producing antibiotic components. Cotton wilt induced by mycelial growth inhibition and toxin production.	Lin <i>et al.</i> , 2013
3.	<i>F. oxysporum</i> f. sp. <i>cubense</i> race 4 <i>Fusarium oxysporum</i> f. sp. <i>cubense</i>	<i>Burkholderia cepacia</i> is a kind of bacteria. Strains 84 and 4B of <i>Pseudomonas putida</i> . Strains of <i>Bacillus cereus</i> , <i>Acromobacter</i> spp., strains of <i>Bacillus flexus</i> <i>Rhizobium</i> spp., W19 <i>Bacillus amyloliquefaciens</i>	Colonize the hyphae and macrospores of the fungal pathogens by inducing mycelial deformities. It has been demonstrated that siderophores and secondary metabolites like surfactin, iturin, and bacillomycin D produce a thick biological layer that prevents pathogen development.	Smith <i>et al.</i> , 2003; Thangavelu and Gopi, 2015
4	<i>Fusarium oxysporum</i>	BECS7, BECS4 and BECL5 <i>Pseudomonas fluorescens</i> (Pf1) <i>Bacillus subtilis</i> (EPCO16 and EPC5), <i>Pseudomonas</i> spp.	Pathogen suppression by hydrolytic enzyme synthesis	Amaresan <i>et al.</i> , 2014
5	<i>F. Avenaciarum</i> <i>F. sambucinum</i> <i>F. oxysporum</i>	<i>Bacillus</i> spp.	<i>In vitro</i> antibiosis	Sturz <i>et al.</i> , 1999
6	<i>C. fagacearum</i>	<i>Pseudomonas denitrificans</i> and <i>P. putida</i>	<i>In vitro</i> antagonism and competitive colonization of microbes	Brooks <i>et al.</i> , 1994



Table 3. Management of various rot causing pathogens by endophytic bacteria

Endophytic Bacteria	Isolated from	Disease	Pathogen	Reference
<i>Actinoplanes missouriensis</i>	Lupin roots	Root rot of lupin	<i>Plectosporium tabacinum</i>	El-Tarabily, 2003
<i>Bacillus amyloliquefaciens</i>	Stems, leaves, and roots of the <i>Eleusine indica</i> (weed)	Stem end rot of pitaya	<i>Alternaria alternata</i>	Trung <i>et al.</i> , 2021
<i>Bacillus subtilis</i> subsp. <i>subtilis</i> and <i>B. amyloliquefaciens</i>	Soybean roots	Charcoal rot of soybean	<i>Macrophomina phaseolina</i>	Torres <i>et al.</i> , 2016
<i>Bacillus megaterium</i> and <i>Enterobacter hormaechei</i> subsp. <i>xiangfangensis</i>	Mangroves and other vascular shrubs	Root rot of bean	<i>Fusarium solani</i>	Mutungi <i>et al.</i> , 2022
<i>Bacillus subtilis</i> and <i>Mesorhizobium ciceri</i>	Nodules of chickpea	Root rot of chickpea	<i>Fusarium solani</i>	Egamberdieva <i>et al.</i> , 2017
<i>Bacillus cereus</i> and <i>Pseudomonas aeruginosa</i>	Rhizome of turmeric	Rhizome rot of turmeric	<i>Pythium aphanidermatum</i>	Vinayarani and Prakash, 2018
<i>Bacillus mycoides</i> isolates BP24 from	Sugar beet leaves	Black pod rot of cacao	<i>Phytophthora capsica</i>	Bargabus <i>et al.</i> 2002; Bargabus <i>et al.</i> , 2004; Melnick <i>et al.</i> , 2008
<i>Bacillus pumilis</i>	Germinating sugar beet seeds			
<i>Bacillus cereus</i>	Potato and tomato plants			
<i>Burkholderia gladioli</i>	Healthy corm of saffron	Corm rot of saffron	<i>Fusarium oxysporum</i>	Ahmad <i>et al.</i> , 2021
<i>Bacillus</i> , <i>Lysinibacillus</i> , and <i>Stenotrophomonas</i>	Tomato plants	Root rot of tomato	<i>Rhizoctonia solani</i>	Sahu <i>et al.</i> , 2019
		Collar rot of tomato	<i>Sclerotium rolfsii</i>	
<i>Pseudomonas viridiflava</i>	Apoplastic fluids attained from canola leaves	Black rot of canola	<i>Xanthomonas campestris</i> pv. <i>Campestris</i>	Romero <i>et al.</i> , 2019
		Stem rot of canola	<i>Sclerotinia sclerotiorum</i>	
<i>Burkholderia cepacia</i> and <i>Pseudomonas aeruginosa</i>	Symptomless oil palm root tissues	Basal stem rot of oil palm	<i>Ganoderma boninense</i>	Sapak <i>et al.</i> , 2008
<i>Paenibacillus polymyxa</i>	Spermosphere of the Styrian oil pumpkin	Fruit rot of Styrian oil pumpkins	<i>Didymella bryoniae</i>	Fürnkranz <i>et al.</i> , 2012

Table 4. Role of bioactive compounds secreted by endophytic bacteria against post-harvest diseases

Endophytic bacteria	Secretion of bioactive compound	Role against post-harvest pathogens	References
<i>Bacillus subtilis</i>	Iturin A, lipopolysaccharide	Antifungal activity	Ek-Ramos <i>et al.</i> , 2019
<i>Bacillus</i> sp.	Surfactin, fengycin	Used against bacterial diseases	Jasim <i>et al.</i> , 2016
<i>Bacillus amyloliquefaciens</i> CEIZ-11	Lipopolysaccharide	Antifungal activity	Zouari <i>et al.</i> , 2016
<i>Bacillus</i> strains and <i>Enterobacter</i>	3-Methylbutan-1-ol	Manage postharvest infection of <i>Botrytis cinerea</i> on tomato fruit, as well as control grey mold during storage and transit	Chaouachi <i>et al.</i> , 2021
<i>Bacillus</i> sp. and <i>Exiguobacterium acetylicum</i>	$\alpha$ -Farnesene	Reduces the postharvest infection of litchi fruit caused by <i>Peronophythora litchii</i>	Zheng <i>et al.</i> , 2019
<i>Bacillus pumilus</i> TM-R	Ethanol	Antifungal activity against post-harvest pathogens	Morita <i>et al.</i> , 2019
<i>Pseudomonas aeruginosa</i>	Phenyltetradeca-2,5-dienoate	Antibacterial activity	Pratiwi <i>et al.</i> , 2017
<i>Pseudomonas donghuensis</i> P482	Dimethyl sulphide, S-methyl thioacetate, methyl thiocyanate, dimethyl trisulphide, 1-undecan and HCN	Against post-harvest losses caused by <i>Rhizoctonia solani</i>	Ossowicki <i>et al.</i> , 2017
<i>Pseudomonas fluorescens</i> strain WR-1	Volatile organic compounds (VOCs)	Both antibacterial and antifungal activity	Raza <i>et al.</i> , 2016
<i>Pseudomonas putida</i> BP25	Volatile organic compounds (VOCs)	Antifungal activities against <i>Phytophthora capsici</i>	Sheoran <i>et al.</i> , 2015
<i>Streptomyces lavendulae</i> SPS-33	2-Methyl-butanol and 3-methyl-1-butanol	Check the infection of <i>Ceratocystis fimbriata</i> causes postharvest losses in sweet potato	Li <i>et al.</i> , 2020

Table 5. Effect of endophytic bacteria against phytopathogenic nematodes (PPN)

Endophytic Bacteria	Crop	Plant Pathogenic Nematode (PPN)	Effect of Endophyte on PPN	Reference
<i>Pantoea agglomerans</i> , <i>Cedecea davisae</i> , <i>Enterobacter intermedius</i> , <i>Pseudomonas putida</i> and <i>Pseudomonas Fluorescens</i>	Tomato	<i>Meloidogyne incognita</i>	As a seed treatment, it reduces nematode infestation.	Munif <i>et al.</i> , 2000
<i>Agrobacterium radiobacter</i> , <i>Bacillus pumilus</i> , <i>B. brevis</i> , <i>B. megaterium</i> , <i>B. mycoides</i> , <i>B. licheniformis</i> , <i>Chryseobacterium balustinum</i> , <i>Cedecea davisae</i> , <i>Cytophaga johnsonae</i> , <i>Lactobacillus paracasei</i> , <i>Micrococcus luteus</i> , <i>Micrococcus halobius</i> , <i>Pseudomonas syringae</i> and <i>Stenotrophomonas maltophilia</i>	Tomato	<i>Meloidogyne incognita</i>	Number of galls and egg masses were reduced.	Mekete <i>et al.</i> , 2009
<i>Pseudomonas</i> spp., <i>Bacillus</i> spp., <i>Methlobacterium</i> spp.	Okra	<i>Meloidogyne incognita</i>	The quantity of adult females, egg masses, eggs per egg mass, and root gall index were all reduced.	Vetrivelkalai <i>et al.</i> , 2010
<i>Rhizobium etli</i>	Tomato	<i>Meloidogyne incognita</i>	35 days after nematode inoculation, the quantity of eggs per female was reduced.	Martinuz <i>et al.</i> , 2013
<i>Pantoea agglomerans</i> , <i>Cedecea davisae</i> , <i>Enterobacter</i> spp., <i>Pseudomonas putida</i>	Tomato	<i>Meloidogyne incognita</i>	When used as a root dip and soil drench, it reduced early root penetration by second stage juvenile along with the reduction in gall formation.	Munif <i>et al.</i> , 2013

<i>Bacillus cereus</i> , <i>Methylobacterium</i> sp., <i>Pseudomonas</i> sp.	Tomato	<i>Meloidogyne incognita</i>	Adult female population, egg masses, eggs per egg mass were all reduced.	Hu <i>et al.</i> , 2017; Vetrivelkalai, 2019
<i>Bacillus subtilis</i> (Talc based)	Banana	<i>Meloidogyne incognita</i> , <i>Pratylenchus coffeae</i> , <i>Radopholus similis</i> , <i>Helicotylenchus multicinctus</i>	Reduced nematode population	Jonathan and Umamaheswari, 2006
<i>Streptomyces</i> sp.	Banana	<i>Meloidogyne javanica</i>	J2s inhibition	Su <i>et al.</i> , 2017
<i>Rhizobium etli</i>	Potato	<i>Meloidogyne incognita</i>	Reduced number of galls on roots.	Hallmann <i>et al.</i> , 2001
<i>Pseudomonas fluorescens</i> , <i>P. putida</i> , <i>P. syxantha</i> , and <i>P. aurantiacea</i>	Potato	<i>Globodera rostochiensis</i>	Growth and multiplication of nematode population was reduced.	Trifonova <i>et al.</i> , 2014
<i>Bacillus carotarum</i> , <i>B. cereus</i> , and <i>Pseudomonas pseudoalcaligenes</i>	Potato	<i>Globodera rostochiensis</i>	J2 mortality increased by 67-97%; Reduces the amount of cysts by 51-65% and J2s by 48-76%	Istifadah <i>et al.</i> , 2018

Studies on endophytic bacteria invading plant roots and inhibiting nematode development are few. For this study, we show several instances of endophytes as biocontrol agents of phytopathogenic nematode in a range of crops and forests, despite the fact that regulatory rules may classify endophytes as bio-stimulants or soil supplements and others as biopesticides (Table 5). Endophytes are a poorly explored group of microorganisms especially bacterial endophyte which are capable of producing bioactive compounds that can be utilized to combat numerous plant pathogens. Endophytic bacteria have been sources of bioactive and volatile compounds and have proven to be useful for

different group of plant pathogens. In both the pre-harvest and post-harvest stages, endophytic bacterial and actinomycete strains have been widely used as BCAs against a variety of plant diseases. Therefore, the potential colonization efficacy of endophytes is a crucial characteristic for disease management. In conclusion this review explained how plants harbor diverse endophytic bacterial strains, colonizing their parts and some of them emitting volatile organic compounds (VOCs) with antifungal and/or plant growth promotion activity. Using these natural symbionts provides a chance to increase crop production while minimizing the use of hazardous pesticides against plant diseases. Finally, given the lack of research

on endophytic diversity, there is a high likelihood of discovering novel and unique bacterial strains from unexplored wild/cultivated plants.

#### CONFLICT OF INTEREST

The authors declare that there is no conflict of interest

#### AUTHOR CONTRIBUTIONS

Conceptualization and writing of manuscript: Seweta Srivastava and Aspak; table making: Kanuri Komala Siva Katyayani and Dipshikha Kaushik; reviewing and editing: Seweta Srivastava and Meenakshi Rana; Figure drawing and Grammar editing: Shubham Kumar and Raghavendra Reddy Manda; Reference setting: Manash Shukla and Vinit Pratap Singh. All authors have read and agreed to the published version of the manuscript.

#### FUNDING

There is no any grant available.

#### ACKNOWLEDGEMENT

The authors would like to extend their sincere appreciation to the Lovely Professional University, Phagwara, India.

#### REFERENCES

- Afzal, I., Shinwari, Z. K., Sikandar, S and Shahzad, S. 2019. Plant beneficial endophytic bacteria: Mechanisms, diversity, host range and genetic determinants. *Microbiological Research*, **221**: 36-49.
- Ahmad, T., Bashir, A., Farooq, S and Riyaz-Ul-Hassan, S. 2021. *Burkholderia gladioli* E39CS3, an endophyte of *Crocus sativus* Linn., induces host resistance against corm-rot caused by *Fusarium oxysporum*. *Journal of Appl. Microbiology*, **132**: 495–508.
- Amaresan, N., Jayakumar, V and Thajuddin, N. 2014. Isolation and characterization of endophytic bacteria associated with chilli (*Capsicum annum*) grown in coastal agricultural ecosystem. *Indian Journal of Biotechnology*, **13**: 247–255.
- Andrews, L. K. 1992 Biological control in the phyllosphere. *Annual Review of Phytopathology*, **30**: 603–635.
- Aravind, R., Eapen, S. J., Kumar, A., Dinu, A and Ramana, K.V. 2010. Screening of endophytic bacteria and evaluation of selected isolates for suppression of burrowing nematode (*Radopholus similis* Thorne) using three varieties of black pepper (*Piper nigrum* L.). *Crop Protection*, **29**: 318–324.
- Arnold, A. E and Lutzoni, F. 2007. Diversity and host range of foliar fungal endophytes: Are tropical leaves biodiversity hot spots. *Ecology*, **88**: 541- 549.
- Aydi-Ben-Abdallah, R., Jabnoun-Khiareddine, H and Daami-Remadi, M. 2020. Fusarium wilt biocontrol and tomato growth stimulation, using endophytic bacteria naturally associated with *Solanum sodomaeum* and *S. bonariense* plants. *Egyptian Journal of Biological Pest Control*, **30**: 113.
- Backman, P. A., Wilson, M and Murphy, J. F. 1997. Bacteria for biological control of plant diseases. In: N. A. Rehcigl & J. E. Rehcigl (Eds.), Environmentally safe approaches to plant disease control (pp. 95–109). Boca Raton: CRC/Lewis Press.
- Bakker, P. A. H. M., Pieterse, C. M. J and Van Loon, L. C. 2007. Induced systemic resistance by fluorescent *Pseudomonas spp.* *Phytopathology*, **97**: 239–243.
- Bargabus, R. L., Zidack, N. K., Sherwood, J. E and Jacobsen, B. J. 2002. Characterization of systemic resistance in sugar beet elicited by a non-pathogenic, phyllosphere colonizing *Bacillus mycoides*, biological control agent. *Physiological and Molecular Plant Pathology*, **61**: 289-298.
- Bargabus, R. L., Zidack, N. K., Sherwood, J. E and Jacobsen, B. J. 2004. Screening for the identification of potential biological control agents that induce systemic acquired resistance in sugar beet. *Biological Control*, **30**: 342-350.
- Beneduzi, A., Ambrosini, A and Passaglia, L. M. P. 2012. Plant growth-promoting rhizobacteria (PGPR): their potential as antagonists and biocontrol agents. *Genetics and Molecular Biology*, **35**: 1044–1051.

- Bhat, A. A., Shakeel, A., Waqar, S., Handoo, Z. A and Khan, A. A. 2023. Microbes vs. Nematodes: Insights into Biocontrol through Antagonistic Organisms to Control Root-Knot Nematodes. *Plants*, **12**: 451.
- Brooks, D. S., Gonzalez, C. F., Appel, D. N and File, T. H. 1994. Evaluation of endophytic bacteria as potential biocontrol agents for oak wilt. *Biological Control*, **4**: 373–381.
- Burrage, S. G and Jeon, J. 2021. Applications of endophytic microbes in agriculture, biotechnology, medicine, and beyond. *Microbiological Research*, **245**: 126691.
- Busby, P. E., Ridout, M and Newcombe, G. 2016. Fungal endophytes: modifiers of plant disease. *Plant Molecular Biology*, **90**: 645e655.
- Castanheira, N. L., Dourado, A. C., Pais, I., Semedo, J., Scotti-Campos, P., Borges, N., Carvalho, G., Barreto Crespo, M. T and Fareleira, P. 2017. Colonization and beneficial effects on annual ryegrass by mixed inoculation with plant growth promoting bacteria. *Microbiology Research*, **198**: 47–55.
- Chaouachi, M., Marzouk, T., Jallouli, S., Elkahoui, S., Gentzittel, L., Ben, C and Djéali, N. 2021. Activity assessment of tomato endophytic bacteria bioactive compounds for the postharvest biocontrol of *Botrytis cinerea*. *Postharvest Biology and Technology*, **172**: 111389.
- Chaudhary, P., Agri, U., Chaudhary, A., Kumar, A and Kumar, G. 2022. Endophytes and their potential in biotic stress management and crop production. *Frontiers in Microbiology*, **13**: 933017.
- Clay, K. 1988. Fungal endophytes of grasses; a defensive mutualism between plants and fungi. *Ecology*, **69**: 10–16.
- Conner, R. L., Hou, A., Balasubramanian, P., McLaren, D. L., Henriquez, M. A., Chang, K.F and McRae, K.B. 2014. Reaction of dry bean cultivars grown in western Canada to root rot inoculation. *Canadian Journal of Plant Science*, **94**:1219-1230.
- Constantin, M. E., de Lamo, F. J., Vlieger, B. V., Rep, M and Takken, F. L. W. 2019. Endophyte-mediated resistance in tomato to *Fusarium oxysporum* is independent of ET, JA, and SA. *Frontiers in Plant Science*, **10**: 979–992.
- Dandurishvili, N., Toklikishvili, N., Ovadis, M., Eliashvili, P., Giorgobiani, N., Keshelava, R., Tediashvili, M., Vainstein, A., Khmel, I., Szegedi, E and Chernin, L. 2011. Broad-range antagonistic rhizobacteria *Pseudomonas fluorescens* and *Serratia plymolithica* suppress *Agrobacterium* crown-gall tumors on tomato plants. *Journal of Applied Microbiology*, **110**: 341–352.
- de Lamo, F. J., Constantin, M. E., Fresno, D. H., Boeren, S., Rep, M and Takken, F. L. W. 2018. Xylem sap proteomics reveals distinct differences between R gene- and endophyte-mediated resistance against *Fusarium* wilt disease in tomato. *Frontiers in Microbiology*, **9**: 2977–2989.
- Degrassi, G and Carpentieri-Pipolo, V. 2020. Bacterial Endophytes Associated to Crops: Novel Practices for Sustainable Agriculture. *Advances in Biochemistry and Biotechnology*, **5**: 1099.
- Dun-chun, H., Jia-sui, Z and Lian-hui, X. 2016. Problems, challenges and future of plant disease management: from an ecological point of view. *Journal of Integrative Agriculture*, **15**(4): 705–715.
- Egamberdieva, D., Wirth, S. J., Shurigin, V. V., Hashem, A and Abd\_Allah, E. F. 2017. Endophytic Bacteria Improve Plant Growth, Symbiotic Performance of Chickpea (*Cicer arietinum* L.) and Induce Suppression of Root Rot Caused by *Fusarium solani* under Salt Stress. *Frontiers in Microbiology*, **8**: 1887.
- Ek-Ramos, M. J., Gomez-Flores, R., Orozco-Flores, A. A., Rodríguez-Padilla, C., González-Ochoa, G and Tamez-Guerra, P. 2019. Bioactive products from plant endophytic gram-positive bacteria. *Frontiers in Microbiology*, **10**: 463.
- El-Tarabily, K. A. 2003. An endophytic chitinase-producing isolate of *Actinoplanes*

- missouriensis*, with potential for biological control of root rot of lupin caused by *Plectosporium tabacinum*. *Australian Journal of Botany*, **51**: 257–266.
- Erwin, D. C and Ribeiro, O. K. 1996. *Phytophthora Diseases Worldwide*. Saint Paul, MN: *American Phytopathological Society (APS Press)*, p.562.
- Estrada, A. E. R., Jonkers, W., Kistler, H. C and May, G. 2012. Interactions between *Fusarium verticillioides*, *Ustilago maydis* and *Zea mays*: An endophyte, a pathogen, and their shared plant host. *Fungal Genetics and Biology*, **49**: 578-587.
- Fadji, A. E., Babalola. O. O. 2020. Elucidating Mechanisms of Endophytes Used in Plant Protection and other Bioactivities with Multifunctional Prospects. *Frontiers in Bioengineering and Biotechnology*, **8**: 467.
- Fisher, P. J and Petrini, O. 1992. Fungal saprobes and pathogens as endophytes of rice (*Oryza sativa* L.). *New Phytologist*, **120**: 137-143.
- Fürnkranz, M., Lukesch, B., Müller, H., Huss, H., Grube, M and Berg, G. 2012. Microbial diversity inside pumpkins: Microhabitat-specific communities display a high antagonistic potential against phytopathogens. *Microbial Ecology*, **63**: 418–428.
- Ganley, R. J., Snieszko, R. A and Newcombe, G. 2008. Endophyte-mediated resistance against white pine blister rust in *Pinus monticola*. *Forest Ecology and Management*, **255**: 2751-2760.
- Gond, S. K., Bergen, M. S., Torres, M. S and White, J. F. Jr. 2015. Endophytic *Bacillus* spp. produce antifungal lipopeptides and induce host defence gene expression in maize. *Microbiology Research*, **172**: 79–87.
- Hacquard, S., Garrido-Oter, R., González, A., Spaepen, S., Ackermann, G., Lebeis, S., McHardy, A. C., Dangl, J. L., Knight, R., Ley, R and Schulze-Lefert, P. 2015. Microbiota and host nutrition across plant and animal kingdoms. *Cell Host & Microbe*. **17**: 603–616.
- Hallman, J., Quadt-Hallmann, A., Mahaffee, W. F. and Klopper, J. W. 1997. Bacterial endophytes in agricultural crops. *Canadian Journal of Microbiology*, **43**: 895–914.
- Hallmann, J., Quadt-Hallmann, A., Miller, W. G., Sikora, R. A and Lindow, S.E. 2001. Endophytic colonization of plants by the biocontrol agent *Rhizobium etli* G12 in relation to *Meloidogyne incognita* infection. *Phytopathology*, **91**: 415–422.
- Harman, G., Khadka, R., Doni, F and Uphoff, N. 2021. Benefits to Plant Health and Productivity from Enhancing Plant Microbial Symbionts. *Frontiers in Plant Science*, **11**: 610065.
- Harni, R., Saefudin, Sasmita, K. D., Sakiroh and Amaria, W. 2023. Efficacy of organic fertilizer, biofertilizer and endophytic bacteria to control nematodes in *robusta coffea*. IOP Conference Series: *Earth and Environmental Science*, **1208**: 012017.
- Hashem, A., Abd Allah, E. F., Alqarawi, A., Al-Huqail, A. A., Wirth, S and Egamberdieva, D. 2016. The interaction between arbuscular mycorrhizal fungi and endophytic bacteria enhances plant growth of *Acacia gerrardii* under salt stress. *Frontiers in Plant Science*, **7**: 1089.
- Hashem, A. H., Attia, M. S., Kandil, E. K., Fawzi, M. M., Abdelrahman, A. S., Khader, M. S., Khodaira, M. A., Emam, A. E., Goma, M. A and Abdelaziz, A. M. 2023. Bioactive compounds and biomedical applications of endophytic fungi: a recent review. *Microbial Cell Factories*, **22**(1): 107.
- Henry, G., Deleu, M., Jourdan, E., Thonart, P and Ongena, M. 2011. The bacterial lipopeptide surfactin targets the lipid fraction of the plant plasma membrane to trigger immune-related defence responses. *Cellular Microbiology*, **13**: 1824–1837.
- Horn, W. S., Simmonds, M. S. J., Schwartz, R. E. and Blaney, W. M. 1995. Phomopsichalasin, a novel antimicrobial agent from an endophytic *Phomopsis* sp. *Tetrahedron*, **51**: 3969–3978.
- Hu, H., Chen, Y., Wang, Y., Tang, Y., Chen, S and Yan, S. 2017. Endophytic *Bacillus cereus*

- effectively controls *Meloidogyne incognita* on tomato plants through rapid rhizosphere occupation and repellent action. *Plant Disease*, **101**: 448–455.
- Huang, X., Ren, J., Li, P., Feng, S., Dong, P and Ren, M. 2021. Potential of microbial endophytes to enhance the resistance to postharvest diseases of fruit and vegetables. *Journal of the Science of Food and Agriculture*, **101**(5): 1744–1757.
- Istifadah, N., Pratama, N., Taqwim, S and Sunarto, T. 2018. Effects of bacterial endophytes from potato roots and tubers on potato cyst nematode (*Globodera rostochiensis*). *Biodiversitas*, **19**: 47–51.
- Jasim, B., Sreelakshmi, K. S., Mathew, J and Radhakrishnan, E. K. 2016. Surfactin, iturin, and fengycin biosynthesis by endophytic *Bacillus* sp. from *Bacopa monnieri*. *Microbial Ecology*, **72**(1): 106–119.
- Jha, P., Kaur, T., Chhabra, I., Panja, A., Paul, S., Kumar, V and Malik, T. 2023. Endophytic fungi: hidden treasure chest of antimicrobial metabolites interrelationship of endophytes and metabolites. *Frontiers in Microbiology*, **14**:1227830.
- Jiang, C. X., Li, J., Zhang, J. M., Jin, X. J., Yu, B., Fang, J and Wu, Q. X. 2019. Isolation, Identification, and Activity Evaluation of Chemical Constituents from Soil Fungus *Fusarium avenaceum* SF-1502 and Endophytic Fungus *Fusarium proliferatum* AF-04. *Journal of Agricultural and Food Chemistry*, **67**: 1839–1846.
- Jonathan, E. I and Umamaheswari, R. 2006. Biomangement of nematodes infesting banana by bacterial endophytes (*Bacillus subtilis*). *Indian Journal of Nematology*, **36**: 6303–6960.
- Kandel, S. L., Joubert, P. M and Doty, S. L. 2017. Bacterial Endophyte Colonization and Distribution within Plants. *Microorganisms*, **5**(4): 77.
- Kang, S. H., Cho, H. S., Cheong, H., Ryu, C. M., Kim, J. F and Park, S. H. 2007. Two bacterial endophytes eliciting boot plant growth promotion and plant defense on pepper (*Capsicum annuum* L.). *Journal of Microbiology and Biotechnology*, **17**: 96–103.
- Kapat, A., Zimand, G and Elad, Y. 1998. Effect of two isolates of *Trichoderma harzianum* on the activity of hydrolytic enzymes produced by *Botrytis cinerea*. *Physiological and Molecular Plant Pathology*, **52**(2): 127–137.
- Kira'ly, L., Barna, B and Kira'ly, Z. 2007. Plant resistance to pathogen infection: forms and mechanisms of innate and acquired resistance. *Journal of Phytopathology*, **155**: 385–396.
- Kirchhof, G., Reis, V. M., Baldani, J. I., Eckert, B., Döbereiner, J and Hartmann, A. 1997. Occurrence, physiological and molecular analysis of endophytic diazotrophic bacteria in gramineous energy plants. *Plant and Soil*, **194**: 45–55.
- Kobayashi, D. Y and Palumbo, J. D. 2000. Bacterial endophytes and their effects on plants and uses in agriculture. In: *Microbial Endophytes*; Bacon, C. W., White, J., Eds.; CRC Press: Boca Raton, FL, USA, pp. 199–233.
- Kumari, M., Qureshi, K. A., Jaremko, M., White, J., Singh, S. K., Sharma, V. K., Singh, K. K., Santoyo, G., Puopolo, G and Kumar, A. 2022. Deciphering the role of endophytic microbiome in postharvest diseases management of fruits: Opportunity areas in commercial up-scale production. *Frontiers in Plant Science*, **13**:1026575.
- Kusari, S., Hertweck, C and Spiteller, M. 2012. Chemical ecology of endophytic fungi: origins of secondary metabolites. *Chemistry & Biology*, **19**: 792–798.
- Lamichhane, J. R., Dürr, C., Schwanck, A. A., Robin, M. H., Sarthou, J. P., Cellier, V., Messéan, A and Aubertot, J. N. 2017. Integrated management of damping-off diseases. *A review. Agronomy for Sustainable Development*, **37**:10.
- Li, X., Li, B., Cai, S., Zhang, Y., Xu, M., Zhang, C., Yuan, B., Xing, K and Qin, S. 2020. Identification of *Rhizospheric actinomycetes*



- Streptomyces lavendulae* SPS-33 and the inhibitory effect of its volatile organic compounds against *Ceratocystis fimbriata* in postharvest sweet potato. *Microorganisms*, **8**(3): 319.
- Lin, T., Zhao, L., Yang, Y., Guan, Q and Gong, M. 2013. Potential of endophytic bacteria isolated from *Sophora alopecuroides* nodule in biological control against *Verticillium wilt* disease. *AJCS*, **7**(1): 139–146.
- Liu, H., Chen, Y., Li, H., Li, S., Tan, H., Liu, Z., Li, D., Liu, H and Zhang, W. 2019. Four new metabolites from the endophytic fungus *Diaporthe lithocarpus* A740. *Fitoterapia*, **137**: 104260.
- Liu, J. Y., Song, Y. C., Zhang Z, *et al.*, 2004. *Aspergillus fumigatus* CY018, an endophytic fungus in *Cynodon dactylon* as a versatile producer of new and bioactive metabolites. *Journal of Biotechnology*, **114**: 279–287.
- Liu, Y., Ponpandian, L. N., Kim, H., Jeon, J., Hwang, B.S., Lee, S.K., Park, SC and Bae, H. (2019) Distribution and diversity of bacterial endophytes from four *Pinus* species and their efficacy as biocontrol agents for devastating pine wood nematodes. *Scientific Reports* **9**: 12461
- Lu H, Zou, W. X., Meng, J. C., *et al.* (2000) New bioactive metabolites produced by *Colletotrichum sp.*, an endophytic fungus in *Artemisia annua*. *Plant Science* **151**: 67–73.
- Lugtenberg, B., Kamilova, F. (2009) Plant-growth-promoting-rhizobacteria. *Annual Review of Microbiology*, **63**: 541–556
- Malfanova, N. V. 2013. Endophytic bacteria with plant growth promoting and biocontrol abilities. Thesis p. 169.
- Malinowski, D. P., Alloush, G. A and Belesky, D. P. 2000. Leaf endophyte *Neotyphodium coenophialum* modifies mineral uptake in tall fescue. *Plant and Soil*, **227**: 115–126.
- Martínez, L., Caballero-Mellado, J., Orozco, J and Martínez-Romero, E. 2003. Diazotrophic bacteria associated with banana (*Musa* spp.). *Plant and Soil*, **257**: 35–47.
- Martinuz, A., Schouten, A and Sikora, R.A. 2013. Post-infection development of *Meloidogyne incognita* on tomato treated with the endophytes *Fusarium oxysporum* strain Fo162 and *Rhizobium etli* strain G12. *Biological Control*, **58**: 95–104.
- Meja, L. C., Rojas, E. I., Maynard, Z., Bael, S. V., Elizabeth Arnold, A., Hebbbar, P., Samuels, G.J., Robbins, N and Herre, E. A. 2008. Endophytic fungi as biocontrol agents of *Theobroma cacao* pathogens. *Biological Control*, **46**: 4-14.
- Mekete, T., Hallmann, J., Kiewnick, S and Sikora, R. 2009. Endophytic bacteria from Ethiopian coffee plants and their potential to antagonize *Meloidogyne incognita*. *Nematology*, **11**: 117–127.
- Melnick, R. L., Zidack, N. K., Bailey, B. A., Maximova, S.N., Gultinan, M and Backman, P.A. 2008. Bacterial endophytes: *Bacillus* spp. from annual crops as potential biological control agents of black pod rot of cacao. *Biological Control*, **46**: 46-56.
- Mercado-Blanco, J., Rodríguez-Jurado, D., Hervás, A and Jiménez-Díaz, R.M. 2004. Suppression of *Verticillium wilt* in olive planting stocks by root-associated fluorescent *Pseudomonas* spp. *Biological Control*, **30**: 474–486.
- Morita, T., Tanaka, I., Ryuda, N., Ikari, M., Ueno, D., Someya, T. 2019 Antifungal spectrum characterization and identification of strong volatile organic compounds produced by *Bacillus pumilus* TM-R. *Heliyon*, **5**: e01817.
- Muhae-ud-Din, G., Moosa, A., Ghummen, U. F., Jabran, M., Abbas, A., Naveed, M., Jabbar, A. and Ali, M. A. 2018. Host status of commonly planted ornamentals to *Meloidogyne incognita* and management through endophytic bacteria. *Pakistan Journal of Zoology*, **50**: 1393–1402.
- Munif, A., Hallmann, J and Sikora, R. A. 2000. Evaluation of the biocontrol activity of endophytic bacteria from tomato against *Meloidogyne incognita*. *Med. Fac. Landbouwkund Universiteit Gent*, **65**: 471–480.

- Munif, A., Hallmann, J and Sikora, R. A. 2013. The influence of endophytic bacteria on *Meloidogyne incognita* infection and tomato plant growth. *Journal of the International Society for Southeast Asian Agricultural Sciences*, **19**: 68–74.
- Muthukumar, A and Bhaskaran, R. 2007. Efficacy of anti-microbial metabolites of *Pseudomonas fluorescens* (Trevisan) Migula. against *Rhizoctonia solani* Khun and *Pythium sp.* *Journal of Biological Control*, **21**: 105–110.
- Mutungi, P. M., Wekesa, V. W., Onguso, J., Kanga, E., Baleba, S. B. S and Boga, H. I. 2022. Culturable Bacterial endophytes associated with shrubs growing along the draw-down zone of lake Bogoria, Kenya: Assessment of antifungal potential against *Fusarium solani* and Induction of Bean Root Rot Protection. *Frontiers in Plant Science*, **12**: 796847.
- Ossowicki, A., Jafra, S and Garbeva, P. 2017. The antimicrobial volatile power of the rhizospheric isolate *Pseudomonas donghuensis* P482. *PLoS One*, **12**: e0174362.
- Padgham, J. L and Sikora, R.A. 2007. Biological control potential and modes of action of *Bacillus megaterium* against *Meloidogyne graminicola* on rice. *Crop Protection*, **26**: 971–977
- Pennell, C., Rolston, M., De Bonth, A., Simpson, W. R and Hume D. E. 2010. Development of a bird-deterrent fungal endophyte in turf tall fescue. *New Zealand Journal of Agricultural Research*, **53**: 145–150.
- Pierson, L. S 3rd and Pierson, E. A. 2010. Metabolism and function of phenazines in bacteria: impacts on the behavior of bacteria in the environment and biotechnological processes. *Applied Microbiology and Biotechnology*, **86**(6):1659-1670.
- Pleban, S., Chernin, L and Chet, I. 1997. Chitinolytic activity of an endophytic strain of *Bacillus cereus*. *Letters in Applied Microbiology*, **25**: 284–288.
- Ponpandian, L. N., Rim, S. O., Shanmugam, G., Jeon, J., Park, Y. H., Lee, S. K and Bae, H. 2019. Phylogenetic characterization of bacterial endophytes from four Pinus species and their nematicidal activity against the pine wood nematode. *Scientific Reports*, **9**: 12457.
- Pratella, G., Mari, M., Guizzardi, F and Folchi, A. 1993. Preliminary studies on the efficiency of endophytes in the biological control of the postharvest pathogens *Monilinia laxa* and *Rhizopus stolonifer* in stone fruit. *Postharvest Biological Technology*, **3**: 361–368.
- Prieto, P., Schilirò, E., Maldonado-González, M. M., Valderrama, R., Barroso-Albarracín, J. B. and Mercado-Blanco, J. 2011. Root hairs play a key role in the endophytic colonization of olive roots by *Pseudomonas spp.* with biocontrol activity. *Microbial Ecology*, **62**: 435–445.
- Rai, M and Agarkar, G. 2016. Plant-fungal interactions: What triggers the fungi to switch among lifestyles? *Critical Review in Microbiology*, **42**(3): 428-438.
- Ramkumar, G., Yu, S. M and Lee, Y. H. 2013. Influence of light qualities on antifungal lipopeptide synthesis in *Bacillus amyloliquefaciens* JBC36. *European Journal of Plant Pathology*, **137**: 243–248.
- Rangjaroen, C., Sungthong, R., Rerkasem, B., Teaumroong, N., Noisangiam, R and Lumyong, S. 2017. Untapped Endophytic Colonization and Plant Growth-Promoting Potential of the Genus *Novosphingobium* to Optimize Rice Cultivation. *Microbes and Environment*, **32**: 84–87.
- Raza, W., Ling, N., Liu, D., Wei, Z., Huang, Q and Shen, Q. (2016). Volatile organic compounds produced by *Pseudomonas fluorescens* WR-1 restrict the growth and virulence traits of *Ralstonia solanacearum*. *Microbiological Research*, **192**: 103-113.
- Romero, F. M., Rossi, F. R., Gárriz, A., Carrasco, P and Ruíz, O. A. 2019. A Bacterial Endophyte from Apoplast Fluids Protects Canola Plants from Different Phytopathogens via Antibiosis and Induction of Host Resistance. *Phytopathology*, **109**: 375–383.

- Rosenblueth, M and Martínez-Romero, E. 2006. Bacterial endophytes and their interactions with hosts. *Molecular Plant-Microbe Interactions*, **19**: 827–837.
- Rybakova, D., Cernava, T., Köberl, M., Liebminger, S., Etemadi, M and Berg, G. 2016. Endophytes-assisted biocontrol: novel insights in ecology and the mode of action of *Paenibacillus*. *Plant Soil*, **405**: 125–140.
- Saeed, Q., Xiukang, W., Haider, F.U., Kučerik, J., Mumtaz, M.Z., Holatko, J., Naseem, M., Kintl, A., Ejaz, M., Naveed, M., Brtnicky, M and Mustafa, A. 2021. Rhizosphere Bacteria in Plant Growth Promotion, Biocontrol, and Bioremediation of Contaminated Sites: A Comprehensive Review of Effects and Mechanisms. *International Journal of Molecular Sciences*, **22**(19): 10529.
- Sahu, P. K., Singh, S., Gupta, A., Singh, U.B., Brahmaprakash, G and Saxena, A. K. 2019. Antagonistic potential of bacterial endophytes and induction of systemic resistance against collar rot pathogen *Sclerotium rolfsii* in tomato. *Biological Control*, **137**: 104014.
- Sapak, Z., Meon, S and Ahmad, Z. A. M. 2008. Effect of endophytic bacteria on growth and suppression of *Ganoderma* infection in oil palm. *International Journal of Agriculture and Biology*, **10**: 127–132.
- Savado, A., Tapi, A., Chollet, M., Wathélet, B., Traore, A. S and Jacques, P. 2011. Identification of surfactin producing strains in Soumbala and Bikalga fermented condiments using polymerase chain reaction and matrix assisted laser desorption /ionization-mass spectrometry methods. *International Journal of Food Microbiology*, **151**: 299–306.
- Schulz, B., Haas, S., Junker, C., Andre'e, N and Schobert, M. 2015. Fungal endophytes are involved in multiple balanced antagonisms. *Current Science*, **109**(1): 39-45.
- Senthilkumar, M., Govindasamy, V and Annapurna, K. 2007. Role of antibiosis in suppression of charcoal rot disease by soybean endophyte *Paenibacillus* sp. HKA-15. *Current Microbiology*, **55**: 25–29.
- Shahzad, R., Khan, A. L., Bilal, S., Asaf, S and Lee, I. J. 2017. Plant growth-promoting endophytic bacteria versus pathogenic infections: an example of *Bacillus amyloliquefaciens* RWL-1 and *Fusarium oxysporum* f. sp. *lycopersici* in tomato. *Peer-reviewed Journal*, **5**: 3107.
- Sharma, H., Rai, A. K., Dahiya, D., Chettri, R and Nigam, P. S. 2021. Exploring endophytes for in vitro synthesis of bioactive compounds similar to metabolites produced in vivo by host plants. *AIMS Microbiology*, **7**(2):175-199.
- Sheoran, N., Nadakkakath, A. V., Munjal, V., Kundu, A., Subaharan, K., Venugopal, V., Rajamma, S., Eapen, S. J and Kumar, A. 2015. Genetic analysis of plant endophytic *Pseudomonas putida* BP25 and chemoprofiling of its antimicrobial volatile organic compounds. *Microbiological Research*, **173**: 66–78.
- Shi, J., Liu, A., Li, X. and Chen, W. 2013. Control of *Phytophthora nicotianae* disease, induction of defense responses and genes expression of papaya fruits treated with *Pseudomonas putida* MGP1. *Journal of the Science of Food and Agriculture*, **93**(3): 568–574.
- Siddiqui, Z.A. and Mahmood, I. 1999. Role of bacteria in the management of plant parasitic nematodes: A review. *Bioresource Technology*, **69**: 167–179.
- Smith, L., Keef, D. O., Smith, M and Hamill, S. 2003. The benefits of applying rhizobacteria to tissue cultured bananas. *Banana Topics Newsletter*, **33**: 1–4.
- Solis, M. J. L., Cruz, T. E. D., Schnittler, M and Unterseher, M. 2016. The diverse community of leaf-inhabiting fungal endophytes from Philippine natural forests reflects phylogenetic patterns of their host plant species *Ficus benjamina*, *F. elastica* and *F. reli-giosa*. *Mycoscience*, **57**: 96e106.
- Sturz, A. V., Christie, B. R., Matheson, B. G., Arsenault, W. J and Buchman, N. A. 1999. Endophytic bacterial communities in the

- periderm of potato tubers and their potential to improve resistance to soil borne plant pathogens. *Plant Pathology*, **48**: 360–369.
- Su, L., Shen, Z., Ruan, Y., Tao, C., Chao, Y., Li, R and Shen, Q. 2017. Isolation of antagonistic endophytes from banana roots against *Meloidogyne javanica* and their effects on soil nematode community. *Frontiers in Microbiology*, **8**: 2070.
- Suryanarayana, T. S., Rajulu, G and Vidal, S. 2016. Biological control through fungal endophytes: Gaps in knowledge hindering success. *Current Biotechnology*, **5**: 1e13.
- Thangavelu, R and Gopi, M. 2015. Field suppression of Fusarium wilt disease in banana by the combined application of native endophytic and rhizospheric bacterial isolates possessing multiple functions. *Phytopathologia Mediterranea*, **54**(2): 241–252.
- Torres, M. J., Pérez Brandan, C., Petroselli, G., Erra-Balsells, R and Audisio, M.C. 2016. Antagonistic effects of *Bacillus Subtilis* subsp. *subtilis* and *B. amyloliquefaciens* against *Macrophomina phaseolina*: SEM study of fungal changes and UV-MALDI-TOF MS analysis of their bioactive compounds. *Microbiological Research*, **182**: 31–39.
- Tran, T. P. H., Wang, S. L., Nguyen, V. B., Tran, D. M., Nguyen, D. S and Nguyen, A. D. 2019. Study of novel endophytic bacteria for biocontrol of black pepper root-knot nematodes in the central highlands of Vietnam. *Agronomy*, **9**: 714.
- Trifonova, Z., Tsvetkov, I., Bogatzevska, N and Batchvarova, R. 2014. Efficiency of *Pseudomonas spp.* for biocontrol of the potato cyst nematode *Globodera rostochiensis* (Woll.). *Bulgarian Journal of Agricultural Science*, **20**: 666–669.
- Trung, D. Q., Anh, L. T., Thuy, N. T., Van, D. M and Hang, T. T. 2021. Endophytic bacteria isolated from a weed plant as a potential biocontrol agent against stem end rot pathogen of pitaya in Vietnam. *Egyptian Journal of Biological Pest Control*, **31**:106.
- Uroz, S., Angelo-Picard, C. D., Carlier, A., Elasri, M., Sicot, C., Petit, A., Oger, P., Faure, D and Dessaux, Y. 2003. Novel bacteria degrading N-acylhomoserine lactones and their use as quenchers of quorum-sensing-regulated functions of plant-pathogenic bacteria. *Microbiology*, **149**: 1981–1989.
- Vetrivelkai, P. 2019. Evaluation of endophytic bacterial isolates against root knot nematode, *Meloidogyne incognita* in tomato under glasshouse condition. *International Journal of Current Microbiology and Applied Sciences*, **8**: 2584–2589.
- Vetrivelkai, P., Sivakumar, M and Jonathan, E. I. 2010. Biocontrol potential of endophytic bacteria on *Meloidogyne incognita* and its effect on plant growth in bhendi. *Journal of Biopesticides*, **3**: 452–457.
- Vinayarani, G., Prakash, H. S. (2018) Growth Promoting Rhizospheric and Endophytic Bacteria from *Curcuma longa* L. as Biocontrol Agents against Rhizome Rot and Leaf Blight Diseases. *Plant Pathology Journal*, **34**(3): 218-235.
- Vitullo, D., Di Pietro, A., Romano, A., Lanzotti, V and Lima, G. 2012. Role of new bacterial surfactins in the antifungal interaction between *Bacillus amyloliquefaciens* and *Fusarium oxysporum*. *Plant Pathology*, **61**(4): 689–699.
- Wang, S. S., Liu, J. M., Sun, J., Sun, Y. F., Liu, J. N., Jia, N., Fan, B and Dai, X.F. 2019. Diversity of culture-independent bacteria and antimicrobial activity of culturable endophytic bacteria isolated from different dendrobium stems. *Scientific Reports*, **9**: 1887.
- Whipps, J. M. 2001. Microbial interactions and biocontrol in the rhizosphere. *Journal of Experimental Botany*, **52**: 487–511.
- Wilson, D. 1995. Endophyte: The evolution of a term, and clarification of its use and definition. *Oikos*, **73**: 274–276.

- Xu, J. X., Li, Z. Y., Lv, X., Yan, H., Zhou, G.Y., Cao, L.X., Yang, Q and He, Y.H. 2020. Isolation and characterization of *Bacillus subtilis* strain 1-L-29, an endophytic-bacteria from *Camellia oleifera* with antimicrobial activity and efficient plant-root colonization. *PLoS ONE*, **15**: e0232096.
- Xu, S. J and Kim, B. S. 2014. Biocontrol of Fusarium Crown and Root Rot and Promotion of Growth of Tomato by Paenibacillus Strains Isolated from Soil. *Mycobiology*, **42**(2):158-166.
- Yadav, A. N., Verma, P., Kour, D., Rana, K. L., Kumar, V., Singh, B., Chauhan, V. S., Sugitha, T., Saxena, A, K and Dhaliwal, H. S. (2017) Plant microbiomes and its beneficial multifunctional plant growth promoting attributes. *International Journal of Environmental Sciences & Natural Resources*, **3**: 1–8
- Yan, J. F., Broughton, S. J., Yang, S. L and Gange, A. C. 2015. Do endophytic fungi grow through their hosts systemically? *Fungal Ecology*, **13**: 53-59.
- Zhang, X., Li, B., Wang, Y., Guo, Q., Lu, X., Li, S and Ma, P. 2013. Lipopeptides, a novel protein, and volatile compounds contribute to the antifungal activity of the biocontrol agent *Bacillus atrophaeus* CAB-1. *Applied Microbiology and Biotechnology*, **97**: 9525–9534.
- Zheng, L., Situ, J. J., Zhu, Q. F., Xi, P. G., Zheng, Y., Liu, H. X., Zhou, X and de Jiang, Z. 2019. Identification of volatile organic compounds for the biocontrol of postharvest litchi fruit pathogen *Peronophythora litchi*. , **155**: 37.
- Zheng, Y. K., Miao, C. P., Chen, H. H., Huang, F. F., Xia, Y. M., Chen, Y. W and Zhao, L. X. 2017. Endophytic fungi harbored in *Panax notoginseng*: Diversity and potential as biological control agents against host plant pathogens of root-rot disease. *Journal of Ginseng Research*, **41**: 353–360.
- Zimand, G., Elad, Y and Chet, I. 1996. Effect of *Trichoderma harzianum* on *Botrytis cinerea* pathogenicity. *Phytopathology*, **86**(11): 1255–1260.
- Zouari, I., Jlaiel, L., Tounsi, S and Trigui, M. 2016. Biocontrol activity of the endophytic *Bacillus amyloliquefaciens* strain CEIZ-11 against *Pythium aphanidermatum* and purification of its bio- active compounds. *Biological Control*, **100**: 54–62.

-----  
**Seweta Srivastava<sup>1\*</sup>, Aspak<sup>1</sup>, Meenakshi Rana<sup>1</sup>, Kanuri Komala Siva Katyayani<sup>2</sup>, Dipshikha Kaushik<sup>2</sup>, Rajeev Kumar<sup>3</sup>, Manash Shukla<sup>1</sup>, Shubham Kumar<sup>1</sup>, Raghavendra Reddy Manda<sup>4</sup> and Vinit Pratap Singh<sup>5</sup>**

<sup>1</sup>School of Agriculture, Lovely Professional University, Phagwara-144 411, Punjab, India

<sup>2</sup>College of Agriculture, Assam Agricultural University, Jorhat-785 013, Assam, India

<sup>3</sup>School of Bioengineering & Biosciences, Lovely Professional University, Phagwara-144 411, Punjab, India

<sup>4</sup>Wageningen University & Research, 6708 PB Wageningen, The Netherlands

<sup>5</sup>College of Agriculture Campus, Kotwa, Azamgarh-276 001, (NDUA&T, Ayodhya), U.P., India

\* Corresponding Author: Seweta Srivastava  
 Email: seweta.21896@lpu.co.in;  
 shalu.bhu2008@gmail.com